

Probabilistic & Unsupervised Learning

Model selection, Hyperparameter optimisation, and Gaussian Processes

Maneesh Sahani


`maneesh@gatsby.ucl.ac.uk`

**Gatsby Computational Neuroscience Unit, and
MSc ML/CSML, Dept Computer Science
University College London**

Term 1, Autumn 2015


Learning model structure

How many clusters in the data?



How smooth should the function be?

Is this input relevant to predicting that output?




What is the order of a dynamical system?

How many states in a hidden Markov model?

SVYDAAAQLTADVKKDLRDSWKVIGSDKKGNG

How many auditory sources in the input?



Model selection

Models (labelled by m) have parameters θ_m that specify the probability of data:

$$P(\mathcal{D}|\theta_m, m).$$

If model is known, learning θ_m means finding posterior or point estimate (ML, MAP, ...).


What if we need to learn the model too?

- ▶ Could combine models into a single “supermodel”, with composite parameter (m, θ_m) .
 - ▶ ML learning will **overfit**: favours most flexible (nested) model with most parameters, even if the data actually come from a simpler one.
 - ▶ Density function on composite parameter space (union of manifolds of different dimensionalities) difficult to define \Rightarrow MAP learning ill-posed.
 - ▶ Joint posterior difficult to compute — dimension of composite parameter varies [but Monte-Carlo methods may sample from much a posterior.]

\Rightarrow Separate model selection step:

$$P(\theta_m, m|\mathcal{D}) = \underbrace{P(\theta_m|m, \mathcal{D})}_{\text{model-specific posterior}} \cdot \underbrace{P(m|\mathcal{D})}_{\text{model selection}}$$

Model complexity and overfitting: a simple example



Model selection

Given models labeled by m with parameters θ_m , identify the “correct” model for data \mathcal{D} .

ML/MAP has no good answer: $P(\mathcal{D}|\theta_m^{\text{ML}})$ is always larger for more complex (nested) models.

Model selection

Given models labeled by m with parameters θ_m , identify the “correct” model for data \mathcal{D} .

ML/MAP has no good answer: $P(\mathcal{D}|\theta_m^{\text{ML}})$ is always larger for more complex (nested) models.

Neyman-Pearson hypothesis testing

- ▶ For **nested** models. Starting with simplest model ($m = 1$), compare (e.g. by likelihood ratio test) **null hypothesis** m to **alternative** $m + 1$. Continue until $m + 1$ is rejected.
- ▶ Usually only valid asymptotically in data number.
- ▶ Conservative (N-P hypothesis tests are asymmetric).

Model selection

Given models labeled by m with parameters θ_m , identify the “correct” model for data \mathcal{D} .

ML/MAP has no good answer: $P(\mathcal{D}|\theta_m^{\text{ML}})$ is always larger for more complex (nested) models.

Neyman-Pearson hypothesis testing

- ▶ For **nested** models. Starting with simplest model ($m = 1$), compare (e.g. by likelihood ratio test) **null hypothesis** m to **alternative** $m + 1$. Continue until $m + 1$ is rejected.
- ▶ Usually only valid asymptotically in data number.
- ▶ Conservative (N-P hypothesis tests are asymmetric).

Likelihood validation

- ▶ Partition data into disjoint *training* and *validation* data sets $\mathcal{D} = \mathcal{D}_{\text{tr}} \cup \mathcal{D}_{\text{vld}}$. Choose model with greatest $P(\mathcal{D}_{\text{vld}}|\theta_m^{\text{ML}})$, with $\theta_m^{\text{ML}} = \text{argmax } P(\mathcal{D}_{\text{tr}}|\theta)$. [Or, better, greatest $P(\mathcal{D}_{\text{vld}}|\mathcal{D}_{\text{tr}}, m)$.]
- ▶ May be biased towards simpler models; often high-variance.
- ▶ **Cross-validation** uses multiple partitions and averages likelihoods.

Model selection

Given models labeled by m with parameters θ_m , identify the “correct” model for data \mathcal{D} .

ML/MAP has no good answer: $P(\mathcal{D}|\theta_m^{\text{ML}})$ is always larger for more complex (nested) models.

Neyman-Pearson hypothesis testing

- ▶ For **nested** models. Starting with simplest model ($m = 1$), compare (e.g. by likelihood ratio test) **null hypothesis** m to **alternative** $m + 1$. Continue until $m + 1$ is rejected.
- ▶ Usually only valid asymptotically in data number.
- ▶ Conservative (N-P hypothesis tests are asymmetric).

Likelihood validation

- ▶ Partition data into disjoint *training* and *validation* data sets $\mathcal{D} = \mathcal{D}_{\text{tr}} \cup \mathcal{D}_{\text{vld}}$. Choose model with greatest $P(\mathcal{D}_{\text{vld}}|\theta_m^{\text{ML}})$, with $\theta_m^{\text{ML}} = \text{argmax } P(\mathcal{D}_{\text{tr}}|\theta)$. [Or, better, greatest $P(\mathcal{D}_{\text{vld}}|\mathcal{D}_{\text{tr}}, m)$.]
- ▶ May be biased towards simpler models; often high-variance.
- ▶ **Cross-validation** uses multiple partitions and averages likelihoods.

Bayesian model selection

- ▶ Choose most likely **model**: $\text{argmax } P(m|\mathcal{D})$.
- ▶ Principled from a probabilistic viewpoint—if true model is in set being considered—but sensitive to assumed priors etc.
- ▶ Can use posterior probabilities to **weight** models for combined predictions (no need to select at all).

Bayesian model selection: some terminology

A **model class** m is a set of distributions parameterised by θ_m , e.g. the set of all possible mixtures of m Gaussians.

The model implies both a **prior** over the parameters $P(\theta_m|m)$, and a **likelihood** of data given parameters (which might require integrating out latent variables) $P(\mathcal{D}|\theta_m, m)$.

The **posterior** distribution over parameters is

$$P(\theta_m|\mathcal{D}, m) = \frac{P(\mathcal{D}|\theta_m, m)P(\theta_m|m)}{P(\mathcal{D}|m)}.$$

The **marginal probability** of the data under model class m is:

$$P(\mathcal{D}|m) = \int_{\Theta_m} P(\mathcal{D}|\theta_m, m)P(\theta_m|m) d\theta_m.$$

(also called the **Bayesian evidence** for model m).

The ratio of two marginal probabilities (or sometimes its log) is known as the **Bayes factor**:

$$\frac{P(\mathcal{D}|m)}{P(\mathcal{D}|m')} = \frac{P(m|\mathcal{D})}{P(m'|\mathcal{D})} \frac{p(m')}{p(m)}$$

The Bayesian Occam's razor

Occam's Razor is a principle of scientific philosophy: of two explanations adequate to explain the same set of observations, the simpler should always be preferred.

Bayesian inference formalises and *automatically* implements a form of Occam's Razor.

Compare model classes m using their posterior probability given the data:


$$P(m|\mathcal{D}) = \frac{P(\mathcal{D}|m)P(m)}{P(\mathcal{D})}, \quad P(\mathcal{D}|m) = \int_{\Theta_m} P(\mathcal{D}|\theta_m, m)P(\theta_m|m) d\theta_m$$

$P(\mathcal{D}|m)$: The probability that *randomly selected* parameter values from the model class would generate data set \mathcal{D} .



Model classes that are **too simple** are unlikely to generate the observed data set.

Model classes that are **too complex** can generate many possible data sets, so again, they are unlikely to generate that particular data set at random.

Like Goldilocks, we favour a model that is **just right**.



Bayesian model comparison: Occam's razor at work



Conjugate-exponential families (recap)

Can we compute $P(\mathcal{D}|m)$?

Conjugate-exponential families (recap)

Can we compute $P(\mathcal{D}|m)$? Sometimes.

Conjugate-exponential families (recap)

Can we compute $P(\mathcal{D}|m)$? Sometimes.

Suppose $P(\mathcal{D}|\boldsymbol{\theta}_m, m)$ is a member of the exponential family:

$$P(\mathcal{D}|\boldsymbol{\theta}_m, m) = \prod_{i=1}^N P(\mathbf{x}_i|\boldsymbol{\theta}_m, m) = \prod_{i=1}^N e^{\mathbf{s}(\mathbf{x}_i)^\top \boldsymbol{\theta}_m - A(\boldsymbol{\theta}_m)}.$$

Conjugate-exponential families (recap)

Can we compute $P(\mathcal{D}|m)$? Sometimes.

Suppose $P(\mathcal{D}|\theta_m, m)$ is a member of the exponential family:

$$P(\mathcal{D}|\theta_m, m) = \prod_{i=1}^N P(\mathbf{x}_i|\theta_m, m) = \prod_{i=1}^N e^{\mathbf{s}(\mathbf{x}_i)^\top \theta_m - A(\theta_m)}.$$

If our prior on θ_m is **conjugate**:

$$P(\theta_m|m) = e^{\mathbf{s}_p^\top \theta_m - n_p A(\theta_m)} / Z(\mathbf{s}_p, n_p)$$

Conjugate-exponential families (recap)

Can we compute $P(\mathcal{D}|m)$? Sometimes.

Suppose $P(\mathcal{D}|\theta_m, m)$ is a member of the exponential family:

$$P(\mathcal{D}|\theta_m, m) = \prod_{i=1}^N P(\mathbf{x}_i|\theta_m, m) = \prod_{i=1}^N e^{\mathbf{s}(\mathbf{x}_i)^\top \theta_m - A(\theta_m)}.$$

If our prior on θ_m is **conjugate**:

$$P(\theta_m|m) = e^{\mathbf{s}_p^\top \theta_m - n_p A(\theta_m)} / Z(\mathbf{s}_p, n_p)$$

then the joint is in the same family:

$$P(\mathcal{D}, \theta_m|m) = e^{(\sum_i \mathbf{s}(\mathbf{x}_i) + \mathbf{s}_p)^\top \theta_m - (N+n_p)A(\theta_m)} / Z(\mathbf{s}_p, p)$$

Conjugate-exponential families (recap)

Can we compute $P(\mathcal{D}|m)$? Sometimes.

Suppose $P(\mathcal{D}|\theta_m, m)$ is a member of the exponential family:

$$P(\mathcal{D}|\theta_m, m) = \prod_{i=1}^N P(\mathbf{x}_i|\theta_m, m) = \prod_{i=1}^N e^{\mathbf{s}(\mathbf{x}_i)^\top \theta_m - A(\theta_m)}.$$

If our prior on θ_m is **conjugate**:

$$P(\theta_m|m) = e^{\mathbf{s}_p^\top \theta_m - n_p A(\theta_m)} / Z(\mathbf{s}_p, n_p)$$

then the joint is in the same family:

$$P(\mathcal{D}, \theta_m|m) = e^{(\sum_i \mathbf{s}(\mathbf{x}_i) + \mathbf{s}_p)^\top \theta_m - (N + n_p) A(\theta_m)} / Z(\mathbf{s}_p, p)$$

and so:

$$P(\mathcal{D}|m) = \int d\theta_m P(\mathcal{D}, \theta_m|m) = Z(\sum_i \mathbf{s}(\mathbf{x}_i) + \mathbf{s}_p, N + n_p) / Z(\mathbf{s}_p, p)$$

Conjugate-exponential families (recap)

Can we compute $P(\mathcal{D}|m)$? Sometimes.

Suppose $P(\mathcal{D}|\theta_m, m)$ is a member of the exponential family:

$$P(\mathcal{D}|\theta_m, m) = \prod_{i=1}^N P(\mathbf{x}_i|\theta_m, m) = \prod_{i=1}^N e^{\mathbf{s}(\mathbf{x}_i)^\top \theta_m - A(\theta_m)}.$$

If our prior on θ_m is **conjugate**:

$$P(\theta_m|m) = e^{\mathbf{s}_p^\top \theta_m - n_p A(\theta_m)} / Z(\mathbf{s}_p, n_p)$$

then the joint is in the same family:

$$P(\mathcal{D}, \theta_m|m) = e^{(\sum_i \mathbf{s}(\mathbf{x}_i) + \mathbf{s}_p)^\top \theta_m - (N + n_p) A(\theta_m)} / Z(\mathbf{s}_p, p)$$

and so:

$$P(\mathcal{D}|m) = \int d\theta_m P(\mathcal{D}, \theta_m|m) = Z(\sum_i \mathbf{s}(\mathbf{x}_i) + \mathbf{s}_p, N + n_p) / Z(\mathbf{s}_p, p)$$

But this is a special case. In general, we need to approximate . . .

Practical Bayesian approaches

- ▶ Laplace approximation:

- ▶ Approximate posterior by a Gaussian centred at the maximum *a posteriori* parameter estimate.

- ▶ Bayesian Information Criterion (BIC)

- ▶ an asymptotic ($N \rightarrow \infty$) approximation.

- ▶ Variational Bayes

- ▶ Lower bound on the marginal probability.
- ▶ Biased estimate.
- ▶ Easy and fast, and often better than Laplace or BIC.

- ▶ Monte Carlo methods:

- ▶ (Annealed) Importance sampling: estimate evidence using samples $\theta^{(i)}$ from arbitrary $f(\theta)$:

$$\sum_i \frac{P(\mathcal{D}|\theta^{(i)}, m)P(\theta^{(i)}|m)}{f(\theta^{(i)})} \rightarrow \int d\theta f(\theta) \frac{P(\mathcal{D}, \theta|m)}{f(\theta)} = P(\mathcal{D}|m)$$

- ▶ “Reversible jump” Markov Chain Monte Carlo: sample from posterior on composite (m, θ_m) .
samples for each $m \propto p(m|\mathcal{D})$.
- ▶ Both exact in the limit of infinite samples, but may have high variance with finite samples.

Not an exhaustive list (Bethe approximations, Expectation propagation, ...)

We will discuss Laplace and BIC now, leaving the rest for the second half of course.

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \boldsymbol{\theta}_m|m) d\boldsymbol{\theta}_m$.

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Idea: approximate $\log P(\mathcal{D}, \theta_m|m)$ to second-order around θ^* .

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Idea: approximate $\log P(\mathcal{D}, \theta_m|m)$ to second-order around θ^* .

$$\int P(\mathcal{D}, \theta_m|m) d\theta_m = \int e^{\log P(\mathcal{D}, \theta_m|m)} d\theta_m$$

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Idea: approximate $\log P(\mathcal{D}, \theta_m|m)$ to second-order around θ^* .

$$\begin{aligned} \int P(\mathcal{D}, \theta_m|m) d\theta_m &= \int e^{\log P(\mathcal{D}, \theta_m|m)} d\theta_m \\ &= \int e^{\log P(\mathcal{D}, \theta_m^*|m) + \nabla \log P(\mathcal{D}, \theta_m^*|m) \cdot (\theta_m - \theta_m^*) + \frac{1}{2} (\theta_m - \theta_m^*)^T \nabla^2 \log P(\mathcal{D}, \theta_m^*|m) (\theta_m - \theta_m^*)} d\theta_m \end{aligned}$$

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Idea: approximate $\log P(\mathcal{D}, \theta_m|m)$ to second-order around θ^* .

$$\begin{aligned}\int P(\mathcal{D}, \theta_m|m) d\theta_m &= \int e^{\log P(\mathcal{D}, \theta_m|m)} d\theta_m \\ &= \int e^{\log P(\mathcal{D}, \theta_m^*|m) + \underbrace{\nabla \log P(\mathcal{D}, \theta_m^*|m)}_{=0} \cdot (\theta_m - \theta_m^*) + \frac{1}{2} (\theta_m - \theta_m^*)^T \underbrace{\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)}_{=-A} (\theta_m - \theta_m^*)} d\theta_m\end{aligned}$$

$A = -\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)$ is the negative Hessian of $\log P(\mathcal{D}, \theta|m)$ evaluated at θ_m^* .

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Idea: approximate $\log P(\mathcal{D}, \theta_m|m)$ to second-order around θ^* .

$$\begin{aligned}\int P(\mathcal{D}, \theta_m|m) d\theta_m &= \int e^{\log P(\mathcal{D}, \theta_m|m)} d\theta_m \\&= \int e^{\log P(\mathcal{D}, \theta_m^*|m) + \underbrace{\nabla \log P(\mathcal{D}, \theta_m^*|m) \cdot (\theta_m - \theta_m^*)}_{=0} + \frac{1}{2} (\theta_m - \theta_m^*)^T \underbrace{\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)}_{=-A} (\theta_m - \theta_m^*)} d\theta_m \\&= \int P(\mathcal{D}, \theta_m^*|m) e^{-\frac{1}{2} (\theta_m - \theta_m^*)^T A (\theta_m - \theta_m^*)} d\theta_m\end{aligned}$$

$A = -\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)$ is the negative Hessian of $\log P(\mathcal{D}, \theta|m)$ evaluated at θ_m^* .

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Idea: approximate $\log P(\mathcal{D}, \theta_m|m)$ to second-order around θ^* .

$$\begin{aligned}\int P(\mathcal{D}, \theta_m|m) d\theta_m &= \int e^{\log P(\mathcal{D}, \theta_m|m)} d\theta_m \\&= \int e^{\log P(\mathcal{D}, \theta_m^*|m) + \underbrace{\nabla \log P(\mathcal{D}, \theta_m^*|m) \cdot (\theta_m - \theta_m^*)}_{=0} + \frac{1}{2} (\theta_m - \theta_m^*)^T \underbrace{\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)}_{=-A} (\theta_m - \theta_m^*)} d\theta_m \\&= \int P(\mathcal{D}, \theta_m^*|m) e^{-\frac{1}{2} (\theta_m - \theta_m^*)^T A (\theta_m - \theta_m^*)} d\theta_m \\&= P(\mathcal{D}|\theta_m^*, m) P(\theta_m^*|m) (2\pi)^{\frac{d}{2}} |A|^{-\frac{1}{2}}\end{aligned}$$

$A = -\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)$ is the negative Hessian of $\log P(\mathcal{D}, \theta|m)$ evaluated at θ_m^* .

Laplace approximation

We want to find $P(\mathcal{D}|m) = \int P(\mathcal{D}, \theta_m|m) d\theta_m$.

As data size N grows (relative to parameter count d), θ_m becomes more constrained
 $\Rightarrow P(\mathcal{D}, \theta_m|m) \propto P(\theta_m|\mathcal{D}, m)$ becomes concentrated on posterior mode θ_m^* .

Idea: approximate $\log P(\mathcal{D}, \theta_m|m)$ to second-order around θ^* .

$$\begin{aligned}\int P(\mathcal{D}, \theta_m|m) d\theta_m &= \int e^{\log P(\mathcal{D}, \theta_m|m)} d\theta_m \\&= \int e^{\log P(\mathcal{D}, \theta_m^*|m) + \underbrace{\nabla \log P(\mathcal{D}, \theta_m^*|m) \cdot (\theta_m - \theta_m^*)}_{=0} + \frac{1}{2} (\theta_m - \theta_m^*)^T \underbrace{\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)}_{=-A} (\theta_m - \theta_m^*)} d\theta_m \\&= \int P(\mathcal{D}, \theta_m^*|m) e^{-\frac{1}{2} (\theta_m - \theta_m^*)^T A (\theta_m - \theta_m^*)} d\theta_m \\&= P(\mathcal{D}|\theta_m^*, m) P(\theta_m^*|m) (2\pi)^{\frac{d}{2}} |A|^{-\frac{1}{2}}\end{aligned}$$

$A = -\nabla^2 \log P(\mathcal{D}, \theta_m^*|m)$ is the negative Hessian of $\log P(\mathcal{D}, \theta|m)$ evaluated at θ_m^* .

This is equivalent to approximating the posterior by a Gaussian: an approximation which is asymptotically correct.

Bayesian Information Criterion (BIC)

BIC can be obtained from the Laplace approximation:

$$\log P(\mathcal{D}|m) \approx \log P(\theta_m^*|m) + \log P(\mathcal{D}|\theta_m^*, m) + \frac{d}{2} \log 2\pi - \frac{1}{2} \log |A|$$

We have

$$A = \nabla^2 \log P(\mathcal{D}, \theta^*|m) = \nabla^2 \log P(\mathcal{D}|\theta^*, m) + \nabla^2 \log P(\theta^*|m)$$

So as the number of iid data $N \rightarrow \infty$, A grows as $NA_0 + \text{constant}$ for a fixed matrix A_0 .
 $\Rightarrow \log |A| \rightarrow \log |NA_0| = \log(N^d |A_0|) = d \log N + \log |A_0|$.

Retaining only terms that grow with N we get:

$$\log P(\mathcal{D}|m) \approx \log P(\mathcal{D}|\theta_m^*, m) - \frac{d}{2} \log N$$


Properties:

- ▶ Quick and easy to compute.
- ▶ Does not depend on prior.
- ▶ We can use the ML estimate of θ instead of the MAP estimate (= as $N \rightarrow \infty$).
- ▶ Related to the “Minimum Description Length” (MDL) criterion.
- ▶ Assumes that in the large sample limit, all the parameters are well-determined (i.e. the model is **identifiable**; otherwise, d should be the number of **well-determined** parameters).
- ▶ Neglects multiple modes (e.g. permutations in a MoG).

Hyperparameters and Evidence optimisation

In some cases, we need to choose between a family of continuously parameterised models.

$$P(\mathcal{D}|\eta) = \int P(\mathcal{D}|\theta)P(\theta|\eta) d\theta$$


hyperparameters

This choice can be made by ascending the gradient in:

- ▶ the exact evidence (if tractable).
- ▶ the approximated evidence (Laplace, EP, Bethe, ...)
- ▶ a free-energy bound on the evidence (Variational Bayes)


or by placing a **hyperprior** on the hyperparameters η , and sampling from the posterior

$$P(\eta|\mathcal{D}) = \frac{P(\mathcal{D}|\eta)P(\eta)}{P(\mathcal{D})}$$

using Markov chain Monte Carlo sampling.

Evidence optimisation in linear regression

Consider simple linear regression:



- Maximize

$$P(y_1 \dots y_N | \mathbf{x}_1 \dots \mathbf{x}_N, C, \sigma^2) = \int P(y_1 \dots y_N | \mathbf{x}_1 \dots \mathbf{x}_N, \mathbf{w}, \sigma^2) P(\mathbf{w} | C) d\mathbf{w}$$

to find optimal values of C, σ .

- Compute the posterior $P(\mathbf{w} | y_1 \dots y_N, \mathbf{x}_1 \dots \mathbf{x}_N, C, \sigma^2)$ given these optimal values.

The evidence for linear regression

- ▶ The posterior on \mathbf{w} is normal: $\Sigma_{\mathbf{w}} = (\frac{XX^T}{\sigma^2} + C^{-1})^{-1}$; $\bar{\mathbf{w}} = \Sigma_{\mathbf{w}} \frac{XY^T}{\sigma^2}$.

Note: X is a matrix where columns are input vectors, and Y is a row vector of corresponding predicted outputs.

The evidence for linear regression

- ▶ The posterior on \mathbf{w} is normal: $\Sigma_{\mathbf{w}} = (\frac{XX^T}{\sigma^2} + C^{-1})^{-1}$; $\bar{\mathbf{w}} = \Sigma_{\mathbf{w}} \frac{XY^T}{\sigma^2}$.

Note: X is a matrix where columns are input vectors, and Y is a row vector of corresponding predicted outputs.

- ▶ The evidence, $\mathcal{E}(C, \sigma^2) = \int P(Y|X, \mathbf{w}, \sigma^2)P(\mathbf{w}|C) d\mathbf{w}$, is given by:

$$\mathcal{E}(C, \sigma^2) = \sqrt{\frac{|2\pi\Sigma_{\mathbf{w}}|}{|2\pi\sigma^2 I| |2\pi C|}} \exp\left(-\frac{1}{2} Y \left(\frac{I}{\sigma^2} - \frac{X^T \Sigma_{\mathbf{w}} X}{\sigma^4}\right) Y^T\right)$$

The evidence for linear regression

- ▶ The posterior on \mathbf{w} is normal: $\Sigma_{\mathbf{w}} = (\frac{XX^T}{\sigma^2} + C^{-1})^{-1}$; $\bar{\mathbf{w}} = \Sigma_{\mathbf{w}} \frac{XY^T}{\sigma^2}$.

Note: X is a matrix where columns are input vectors, and Y is a row vector of corresponding predicted outputs.

- ▶ The evidence, $\mathcal{E}(C, \sigma^2) = \int P(Y|X, \mathbf{w}, \sigma^2)P(\mathbf{w}|C) d\mathbf{w}$, is given by:

$$\mathcal{E}(C, \sigma^2) = \sqrt{\frac{|2\pi\Sigma_{\mathbf{w}}|}{|2\pi\sigma^2 I| |2\pi C|}} \exp\left(-\frac{1}{2} Y \left(\frac{I}{\sigma^2} - \frac{X^T \Sigma_{\mathbf{w}} X}{\sigma^4} \right) Y^T\right)$$

- ▶ For optimization, general forms for the gradients are available. If θ is a parameter in C :

$$\begin{aligned}\frac{\partial}{\partial \theta} \log \mathcal{E}(C, \sigma^2) &= \frac{1}{2} \text{Tr} \left[(C - \Sigma_{\mathbf{w}} - \bar{\mathbf{w}}\bar{\mathbf{w}}^T) \frac{\partial}{\partial \theta} C^{-1} \right] \\ \frac{\partial}{\partial \sigma^2} \log \mathcal{E}(C, \sigma^2) &= \frac{1}{\sigma^2} \left(-N + \text{Tr} [I - \Sigma_{\mathbf{w}} C^{-1}] + \frac{1}{\sigma^2} (Y - \bar{\mathbf{w}}^T X)(Y - \bar{\mathbf{w}}^T X)^T \right)\end{aligned}$$

Automatic Relevance Determination

The most common form of evidence optimization for regression (due to MacKay and Neal) takes $C^{-1} = \text{diag}(\alpha)$ (i.e. $w_i \sim \mathcal{N}(0, \alpha_i^{-1})$) and then optimizes the precisions $\{\alpha_i\}$.

Setting the gradients to 0 and solving gives

$$\alpha_i^{\text{new}} = \frac{1 - \alpha_i [\Sigma_{\mathbf{w}}]_{ii}}{\bar{\mathbf{w}}_i^2}$$
$$(\sigma^2)^{\text{new}} = \frac{(Y - \bar{\mathbf{w}}^T X)(Y - \bar{\mathbf{w}}^T X)^T}{N - \sum_i (1 - [\Sigma_{\mathbf{w}}]_{ii} \alpha_i)}$$


During optimization the α_i s meet one of two fates

| | | | |
|-------------------------------|---------------|---|------------------------|
| $\alpha_i \rightarrow \infty$ | \Rightarrow | $w_i = 0$ | irrelevant input x_i |
| α_i finite | \Rightarrow | $w_i = \text{argmax } P(w_i \mid X, Y, \alpha_i)$ | relevant input x_i |

This procedure, [Automatic Relevance Determination](#) (ARD), yields [sparse](#) solutions that improve on ML regression. (cf. L_1 -regression or LASSO).

Evidence optimisation is also called [maximum marginal likelihood](#) or [ML-2](#) (Type 2 maximum likelihood).

Prediction averaging




Linear regression predicts output y given input vector \mathbf{x} by:

$$y \sim \mathcal{N}(\mathbf{w}^T \mathbf{x}, \sigma^2)$$

Posterior over \mathbf{w} is Gaussian with covariance $\Sigma_{\mathbf{w}} = (\frac{1}{\sigma^2} X X^T + \frac{1}{\tau^2} I)^{-1}$ and mean $\bar{\mathbf{w}} = \frac{1}{\sigma^2} \Sigma_{\mathbf{w}} X Y^T$ (where X is matrix with columns being input vectors, Y is row vector of outputs).

Prediction averaging



Linear regression predicts output y given input vector \mathbf{x} by:

$$y \sim \mathcal{N}(\mathbf{w}^T \mathbf{x}, \sigma^2)$$


Posterior over \mathbf{w} is Gaussian with covariance $\Sigma_{\mathbf{w}} = (\frac{1}{\sigma^2} X X^T + \frac{1}{\tau^2} I)^{-1}$ and mean $\bar{\mathbf{w}} = \frac{1}{\sigma^2} \Sigma_{\mathbf{w}} X Y^T$ (where X is matrix with columns being input vectors, Y is row vector of outputs).

Given a new input vector \mathbf{x}' , the predicted output y' is (integrating out \mathbf{w}):

$$y' | \mathbf{x}' \sim \mathcal{N}(\bar{\mathbf{w}}^T \mathbf{x}', \mathbf{x}'^T \Sigma_{\mathbf{w}} \mathbf{x}' + \sigma^2)$$


the additional variance term $\mathbf{x}'^T \Sigma_{\mathbf{w}} \mathbf{x}'$ comes from the posterior uncertainty in \mathbf{w} .

Marginalised linear regression




Integrate out \mathbf{w}

Marginalised linear regression



Integrate out \mathbf{w} : the joint distribution of y_1, \dots, y_N given $\mathbf{x}_1, \dots, \mathbf{x}_N$ is Gaussian.


Marginalised linear regression



Integrate out \mathbf{w} : the joint distribution of y_1, \dots, y_N given $\mathbf{x}_1, \dots, \mathbf{x}_N$ is Gaussian.

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} \Bigg| \mathbf{x}_1, \dots, \mathbf{x}_N \sim \mathcal{N} \left(\begin{bmatrix} \cdot \\ \cdot \\ \cdot \end{bmatrix}, \begin{bmatrix} \cdot \\ \cdot \\ \cdot \end{bmatrix} \right)$$

Marginalised linear regression



Integrate out \mathbf{w} : the joint distribution of y_1, \dots, y_N given $\mathbf{x}_1, \dots, \mathbf{x}_N$ is Gaussian.
The means and covariances are:


$$E[y_i] =$$

$$E[(y_i - \bar{y}_i)^2] =$$

$$E[(y_i - \bar{y}_i)(y_j - \bar{y}_j)] =$$

$$\left[\begin{array}{c} y_1 \\ y_2 \\ \vdots \\ y_N \end{array} \right] \middle| \mathbf{x}_1, \dots, \mathbf{x}_N \sim \mathcal{N} \left(\left[\begin{array}{c} \vdots \end{array} \right], \left[\begin{array}{c} \vdots \end{array} \right] \right)$$

Marginalised linear regression



Integrate out \mathbf{w} : the joint distribution of y_1, \dots, y_N given $\mathbf{x}_1, \dots, \mathbf{x}_N$ is Gaussian.
The means and covariances are:


$$E[y_i] = E[\mathbf{w}^T \mathbf{x}_i] = \mathbf{0}^T \mathbf{x}_i = 0$$

$$E[(y_i - \bar{y}_i)^2] =$$

$$E[(y_i - \bar{y}_i)(y_j - \bar{y}_j)] =$$

$$\left[\begin{array}{c} y_1 \\ y_2 \\ \vdots \\ y_N \end{array} \right] \Bigg| \mathbf{x}_1, \dots, \mathbf{x}_N \sim \mathcal{N} \left(\left[\begin{array}{c} 0 \\ 0 \\ \vdots \\ 0 \end{array} \right], \left[\begin{array}{c} \end{array} \right] \right)$$

Marginalised linear regression



Integrate out \mathbf{w} : the joint distribution of y_1, \dots, y_N given $\mathbf{x}_1, \dots, \mathbf{x}_N$ is Gaussian.
The means and covariances are:


$$E[y_i] = E[\mathbf{w}^T \mathbf{x}_i] = \mathbf{0}^T \mathbf{x}_i = 0$$

$$E[(y_i - \bar{y}_i)^2] = E[(\mathbf{x}_i^T \mathbf{w})(\mathbf{w}^T \mathbf{x}_i)] + \sigma^2 = \tau^2 \mathbf{x}_i^T \mathbf{x}_i + \sigma^2$$

$$E[(y_i - \bar{y}_i)(y_j - \bar{y}_j)] =$$

$$\left[\begin{array}{c} y_1 \\ y_2 \\ \vdots \\ y_N \end{array} \right] \middle| \mathbf{x}_1, \dots, \mathbf{x}_N \sim \mathcal{N} \left(\left[\begin{array}{c} 0 \\ 0 \\ \vdots \\ 0 \end{array} \right], \left[\begin{array}{ccc} \tau^2 \mathbf{x}_1^T \mathbf{x}_1 + \sigma^2 & & \\ & \tau^2 \mathbf{x}_2^T \mathbf{x}_2 + \sigma^2 & \\ & & \ddots \\ & & & \tau^2 \mathbf{x}_N^T \mathbf{x}_N + \sigma^2 \end{array} \right] \right)$$

Marginalised linear regression



Integrate out \mathbf{w} : the joint distribution of y_1, \dots, y_N given $\mathbf{x}_1, \dots, \mathbf{x}_N$ is Gaussian.

The means and covariances are:


$$E[y_i] = E[\mathbf{w}^T \mathbf{x}_i] = \mathbf{0}^T \mathbf{x}_i = 0$$

$$E[(y_i - \bar{y}_i)^2] = E[(\mathbf{x}_i^T \mathbf{w})(\mathbf{w}^T \mathbf{x}_i)] + \sigma^2 = \tau^2 \mathbf{x}_i^T \mathbf{x}_i + \sigma^2$$

$$E[(y_i - \bar{y}_i)(y_j - \bar{y}_j)] = E[(\mathbf{x}_i^T \mathbf{w})(\mathbf{w}^T \mathbf{x}_j)] = \tau^2 \mathbf{x}_i^T \mathbf{x}_j$$

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} \middle| \mathbf{x}_1, \dots, \mathbf{x}_N \sim \mathcal{N} \left(\begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 \mathbf{x}_1^T \mathbf{x}_1 + \sigma^2 & \tau^2 \mathbf{x}_1^T \mathbf{x}_2 & \cdots & \tau^2 \mathbf{x}_1^T \mathbf{x}_N \\ \tau^2 \mathbf{x}_2^T \mathbf{x}_1 & \tau^2 \mathbf{x}_2^T \mathbf{x}_2 + \sigma^2 & & \tau^2 \mathbf{x}_2^T \mathbf{x}_N \\ \vdots & & \ddots & \vdots \\ \tau^2 \mathbf{x}_N^T \mathbf{x}_1 & \tau^2 \mathbf{x}_N^T \mathbf{x}_2 & \cdots & \tau^2 \mathbf{x}_N^T \mathbf{x}_N + \sigma^2 \end{bmatrix} \right)$$

Marginalised linear regression



$$\mathbf{y}^T \sim \mathcal{N}(\mathbf{0}, \tau^2 \mathbf{X}^T \mathbf{X} + \sigma^2 \mathbf{I})$$

Integrate out \mathbf{w} : the joint distribution of y_1, \dots, y_N given $\mathbf{x}_1, \dots, \mathbf{x}_N$ is Gaussian.

The means and covariances are:

$$E[y_i] = E[\mathbf{w}^T \mathbf{x}_i] = \mathbf{0}^T \mathbf{x}_i = 0$$

$$E[(y_i - \bar{y}_i)^2] = E[(\mathbf{x}_i^T \mathbf{w})(\mathbf{w}^T \mathbf{x}_i)] + \sigma^2 = \tau^2 \mathbf{x}_i^T \mathbf{x}_i + \sigma^2$$

$$E[(y_i - \bar{y}_i)(y_j - \bar{y}_j)] = E[(\mathbf{x}_i^T \mathbf{w})(\mathbf{w}^T \mathbf{x}_j)] = \tau^2 \mathbf{x}_i^T \mathbf{x}_j$$

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} \middle| \mathbf{x}_1, \dots, \mathbf{x}_N \sim \mathcal{N} \left(\begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 \mathbf{x}_1^T \mathbf{x}_1 + \sigma^2 & \tau^2 \mathbf{x}_1^T \mathbf{x}_2 & \cdots & \tau^2 \mathbf{x}_1^T \mathbf{x}_N \\ \tau^2 \mathbf{x}_2^T \mathbf{x}_1 & \tau^2 \mathbf{x}_2^T \mathbf{x}_2 + \sigma^2 & & \tau^2 \mathbf{x}_2^T \mathbf{x}_N \\ \vdots & & \ddots & \vdots \\ \tau^2 \mathbf{x}_N^T \mathbf{x}_1 & \tau^2 \mathbf{x}_N^T \mathbf{x}_2 & \cdots & \tau^2 \mathbf{x}_N^T \mathbf{x}_N + \sigma^2 \end{bmatrix} \right)$$

Predictions with marginalised regression

Now, include the test input vector \mathbf{x}' and test output y' :

$$\begin{bmatrix} Y^T \\ y' \end{bmatrix} \Big| X, \mathbf{x}' \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 X^T X + \sigma^2 I & \tau^2 X^T \mathbf{x}' \\ \tau^2 \mathbf{x}'^T X & \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 \end{bmatrix} \right)$$

We can find $y' | Y$ by the standard multivariate Gaussian result:

$$\begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \textcolor{red}{A} & \textcolor{blue}{C} \\ \textcolor{blue}{C}^T & \textcolor{green}{B} \end{bmatrix} \right) \Rightarrow \mathbf{b} | \mathbf{a} \sim \mathcal{N} \left(\textcolor{blue}{C}^T \textcolor{red}{A}^{-1} \mathbf{a}, \textcolor{green}{B} - \textcolor{blue}{C}^T \textcolor{red}{A}^{-1} \textcolor{blue}{C} \right)$$

Predictions with marginalised regression

Now, include the test input vector \mathbf{x}' and test output y' :

$$\begin{bmatrix} Y^T \\ y' \end{bmatrix} \Big| X, \mathbf{x}' \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 X^T X + \sigma^2 I & \tau^2 X^T \mathbf{x}' \\ \tau^2 \mathbf{x}'^T X & \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 \end{bmatrix} \right)$$

We can find $y' | Y$ by the standard multivariate Gaussian result:

$$\begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} A & C \\ C^T & B \end{bmatrix} \right) \Rightarrow \mathbf{b} | \mathbf{a} \sim \mathcal{N} \left(C^T A^{-1} \mathbf{a}, B - C^T A^{-1} C \right)$$

So

$$\begin{aligned} y' | Y, X, \mathbf{x}' \\ \sim \mathcal{N} \left(\tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} Y^T, \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 - \tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} \tau^2 X^T \mathbf{x}' \right) \end{aligned}$$

Predictions with marginalised regression

Now, include the test input vector \mathbf{x}' and test output y' :

$$\begin{bmatrix} Y^T \\ y' \end{bmatrix} \Big| X, \mathbf{x}' \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 X^T X + \sigma^2 I & \tau^2 X^T \mathbf{x}' \\ \tau^2 \mathbf{x}'^T X & \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 \end{bmatrix} \right)$$

We can find $y' | Y$ by the standard multivariate Gaussian result:

$$\begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} A & C \\ C^T & B \end{bmatrix} \right) \Rightarrow \mathbf{b} | \mathbf{a} \sim \mathcal{N} \left(C^T A^{-1} \mathbf{a}, B - C^T A^{-1} C \right)$$

So

$$y' | Y, X, \mathbf{x}'$$

$$\begin{aligned} &\sim \mathcal{N} \left(\tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} Y^T, \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 - \tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} \tau^2 X^T \mathbf{x}' \right) \\ &\sim \mathcal{N} \left(\frac{1}{\sigma^2} \mathbf{x}'^T \Sigma X Y^T, \mathbf{x}'^T \Sigma \mathbf{x}' + \sigma^2 \right) \quad \Sigma = \left(\frac{1}{\sigma^2} X X^T + \frac{1}{\tau^2} I \right)^{-1} \end{aligned}$$

Predictions with marginalised regression

Now, include the test input vector \mathbf{x}' and test output y' :

$$\begin{bmatrix} Y^T \\ y' \end{bmatrix} \Big| X, \mathbf{x}' \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 X^T X + \sigma^2 I & \tau^2 X^T \mathbf{x}' \\ \tau^2 \mathbf{x}'^T X & \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 \end{bmatrix} \right)$$

We can find $y' | Y$ by the standard multivariate Gaussian result:

$$\begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} A & C \\ C^T & B \end{bmatrix} \right) \Rightarrow \mathbf{b} | \mathbf{a} \sim \mathcal{N} \left(C^T A^{-1} \mathbf{a}, B - C^T A^{-1} C \right)$$

So

$$\begin{aligned} y' | Y, X, \mathbf{x}' &\sim \mathcal{N} \left(\tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} Y^T, \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 - \tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} \tau^2 X^T \mathbf{x}' \right) \\ &\sim \mathcal{N} \left(\frac{1}{\sigma^2} \mathbf{x}'^T \Sigma X Y^T, \mathbf{x}'^T \Sigma \mathbf{x}' + \sigma^2 \right) \quad \Sigma = \left(\frac{1}{\sigma^2} X X^T + \frac{1}{\tau^2} I \right)^{-1} \end{aligned}$$

- Same answer as obtained by integrating wrt **posterior** over \mathbf{w} .

Predictions with marginalised regression

Now, include the test input vector \mathbf{x}' and test output y' :

$$\begin{bmatrix} Y^T \\ y' \end{bmatrix} \Big| X, \mathbf{x}' \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 X^T X + \sigma^2 I & \tau^2 X^T \mathbf{x}' \\ \tau^2 \mathbf{x}'^T X & \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 \end{bmatrix} \right)$$

We can find $y' | Y$ by the standard multivariate Gaussian result:

$$\begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} A & C \\ C^T & B \end{bmatrix} \right) \Rightarrow \mathbf{b} | \mathbf{a} \sim \mathcal{N} \left(C^T A^{-1} \mathbf{a}, B - C^T A^{-1} C \right)$$

So

$$\begin{aligned} y' | Y, X, \mathbf{x}' &\sim \mathcal{N} \left(\tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} Y^T, \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 - \tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} \tau^2 X^T \mathbf{x}' \right) \\ &\sim \mathcal{N} \left(\frac{1}{\sigma^2} \mathbf{x}'^T \Sigma X Y^T, \mathbf{x}'^T \Sigma \mathbf{x}' + \sigma^2 \right) \quad \Sigma = \left(\frac{1}{\sigma^2} X X^T + \frac{1}{\tau^2} I \right)^{-1} \end{aligned}$$

- ▶ Same answer as obtained by integrating wrt **posterior** over \mathbf{w} .
- ▶ Evidence $P(Y|X)$ is just probability under joint Gaussian; also reduces to expression found previously.

Predictions with marginalised regression

Now, include the test input vector \mathbf{x}' and test output y' :

$$\begin{bmatrix} Y^T \\ y' \end{bmatrix} \Big| X, \mathbf{x}' \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \tau^2 X^T X + \sigma^2 I & \tau^2 X^T \mathbf{x}' \\ \tau^2 \mathbf{x}'^T X & \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 \end{bmatrix} \right)$$

We can find $y' | Y$ by the standard multivariate Gaussian result:


$$\begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} A & C \\ C^T & B \end{bmatrix} \right) \Rightarrow \mathbf{b} | \mathbf{a} \sim \mathcal{N} \left(C^T A^{-1} \mathbf{a}, B - C^T A^{-1} C \right)$$

So

$$\begin{aligned} y' | Y, X, \mathbf{x}' &\sim \mathcal{N} \left(\tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} Y^T, \tau^2 \mathbf{x}'^T \mathbf{x}' + \sigma^2 - \tau^2 \mathbf{x}'^T X (\tau^2 X^T X + \sigma^2 I)^{-1} \tau^2 X^T \mathbf{x}' \right) \\ &\sim \mathcal{N} \left(\frac{1}{\sigma^2} \mathbf{x}'^T \Sigma X Y^T, \mathbf{x}'^T \Sigma \mathbf{x}' + \sigma^2 \right) \quad \Sigma = \left(\frac{1}{\sigma^2} X X^T + \frac{1}{\tau^2} I \right)^{-1} \end{aligned}$$


- ▶ Same answer as obtained by integrating wrt **posterior** over \mathbf{w} .
- ▶ Evidence $P(Y|X)$ is just probability under joint Gaussian; also reduces to expression found previously.
- ▶ Thus, Bayesian linear regression can be derived from a **joint, parameter-free** distribution on all the outputs conditioned on all the inputs.

Nonlinear regression



We can also introduce a nonlinear mapping $\mathbf{x} \mapsto \phi(\mathbf{x})$? Each element of $\phi(\mathbf{x})$ is a (nonlinear) feature extracted from \mathbf{x} . May be many more features than elements on \mathbf{x} .

Nonlinear regression




We can also introduce a nonlinear mapping $\mathbf{x} \mapsto \phi(\mathbf{x})$? Each element of $\phi(\mathbf{x})$ is a (nonlinear) feature extracted from \mathbf{x} . May be many more features than elements on \mathbf{x} .

The regression function $f(\mathbf{x}) = \mathbf{w}^\top \phi(\mathbf{x})$ is nonlinear, but outputs Y still jointly Gaussian!

$$Y^\top | X \sim \mathcal{N}(0_N, \tau^2 \Phi^\top \Phi + \sigma^2 I_N)$$

where the i^{th} column of matrix Φ is $\phi(\mathbf{x}_i)$.

Nonlinear regression



We can also introduce a nonlinear mapping $\mathbf{x} \mapsto \phi(\mathbf{x})$? Each element of $\phi(\mathbf{x})$ is a (nonlinear) feature extracted from \mathbf{x} . May be many more features than elements on \mathbf{x} .

The regression function $f(\mathbf{x}) = \mathbf{w}^T \phi(\mathbf{x})$ is nonlinear, but outputs Y still jointly Gaussian!

$$Y^T | X \sim \mathcal{N}(0_N, \tau^2 \Phi^T \Phi + \sigma^2 I_N)$$

where the i^{th} column of matrix Φ is $\phi(\mathbf{x}_i)$.

Proceeding as before, the predictive distribution over y' for a test input \mathbf{x}' is:

$$y' | \mathbf{x}', Y, X \sim \mathcal{N} \left(\tau^2 \phi(\mathbf{x}')^T \Phi K^{-1} Y^T, \tau^2 \phi(\mathbf{x}')^T \phi(\mathbf{x}') + \sigma^2 - \tau^4 \phi(\mathbf{x}')^T \Phi K^{-1} \Phi^T \phi(\mathbf{x}') \right)$$

$$K = \tau^2 \Phi^T \Phi + \sigma^2 I$$

The covariance kernel

$$Y^T|X \sim \mathcal{N}\left(\mathbf{0}_N, \tau^2 \Phi^T \Phi + \sigma^2 I_N\right)$$

The covariance of the output vector Y plays a central role in the development of the theory of Gaussian processes.

The covariance kernel

$$Y^T|X \sim \mathcal{N}\left(\mathbf{0}_N, \tau^2 \Phi^T \Phi + \sigma^2 I_N\right)$$

The covariance of the output vector Y plays a central role in the development of the theory of Gaussian processes.

Define the **covariance kernel** function $K : \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}$ such that if $\mathbf{x}, \mathbf{x}' \in \mathbb{X}$ are two input vectors with corresponding outputs y, y' , then

$$K(\mathbf{x}, \mathbf{x}') = \text{Cov}[y, y'] = E[yy'] - E[y]E[y']$$

In the nonlinear regression example we have $K(\mathbf{x}, \mathbf{x}') = \tau^2 \phi(\mathbf{x})^T \phi(\mathbf{x}') + \sigma^2 \delta_{\mathbf{x}=\mathbf{x}'}$.

The covariance kernel

$$Y^T | X \sim \mathcal{N}(\mathbf{0}_N, \tau^2 \Phi^T \Phi + \sigma^2 I_N)$$

The covariance of the output vector Y plays a central role in the development of the theory of Gaussian processes.

Define the **covariance kernel** function $K : \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}$ such that if $\mathbf{x}, \mathbf{x}' \in \mathbb{X}$ are two input vectors with corresponding outputs y, y' , then

$$K(\mathbf{x}, \mathbf{x}') = \text{Cov}[y, y'] = E[yy'] - E[y]E[y']$$

In the nonlinear regression example we have $K(\mathbf{x}, \mathbf{x}') = \tau^2 \phi(\mathbf{x})^T \phi(\mathbf{x}') + \sigma^2 \delta_{\mathbf{x}=\mathbf{x}'}$.

The covariance kernel has two properties:

- ▶ **Symmetric**: $K(\mathbf{x}, \mathbf{x}') = K(\mathbf{x}', \mathbf{x})$ for all \mathbf{x}, \mathbf{x}' .
- ▶ **Positive semidefinite**: the matrix $[K(\mathbf{x}_i, \mathbf{x}_j)]$ formed by any finite set of input vectors $\mathbf{x}_1, \dots, \mathbf{x}_N$ is positive semidefinite.

The covariance kernel

$$Y^\top | X \sim \mathcal{N}(\mathbf{0}_N, \tau^2 \Phi^\top \Phi + \sigma^2 I_N)$$

The covariance of the output vector Y plays a central role in the development of the theory of Gaussian processes.

Define the **covariance kernel** function $K : \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}$ such that if $\mathbf{x}, \mathbf{x}' \in \mathbb{X}$ are two input vectors with corresponding outputs y, y' , then

$$K(\mathbf{x}, \mathbf{x}') = \text{Cov}[y, y'] = E[yy'] - E[y]E[y']$$

In the nonlinear regression example we have $K(\mathbf{x}, \mathbf{x}') = \tau^2 \phi(\mathbf{x})^\top \phi(\mathbf{x}') + \sigma^2 \delta_{\mathbf{x}=\mathbf{x}'}$.

The covariance kernel has two properties:

- ▶ **Symmetric**: $K(\mathbf{x}, \mathbf{x}') = K(\mathbf{x}', \mathbf{x})$ for all \mathbf{x}, \mathbf{x}' .
- ▶ **Positive semidefinite**: the matrix $[K(\mathbf{x}_i, \mathbf{x}_j)]$ formed by any finite set of input vectors $\mathbf{x}_1, \dots, \mathbf{x}_N$ is positive semidefinite.

Theorem: A covariance kernel $K : \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}$ is symmetric and positive semidefinite if and only if there is a feature map $\phi : \mathbb{X} \rightarrow \mathbb{H}$ such that

$$K(\mathbf{x}, \mathbf{x}') = \phi(\mathbf{x})^\top \phi(\mathbf{x}')$$

The feature space \mathbb{H} can potentially be infinite dimensional.

Regression using the covariance kernel

For non-linear regression, all operations depended on $K(\mathbf{x}, \mathbf{x}')$ rather than explicitly on $\phi(\mathbf{x})$.

Regression using the covariance kernel

For non-linear regression, all operations depended on $K(\mathbf{x}, \mathbf{x}')$ rather than explicitly on $\phi(\mathbf{x})$.

So we can define the joint in terms of K *implicitly* using a (potentially infinite-dimensional) feature map $\phi(\mathbf{x})$.

$$Y|X, K \sim \mathcal{N}(0_N, K(X, X))$$

where the i, j entry in the covariance matrix $K(X, X)$ is $K(\mathbf{x}_i, \mathbf{x}_j)$.

This is called the **kernel trick**.

Regression using the covariance kernel

For non-linear regression, all operations depended on $K(\mathbf{x}, \mathbf{x}')$ rather than explicitly on $\phi(\mathbf{x})$.

So we can define the joint in terms of K *implicitly* using a (potentially infinite-dimensional) feature map $\phi(\mathbf{x})$.

$$Y|X, K \sim \mathcal{N}(0_N, K(X, X))$$

where the i, j entry in the covariance matrix $K(X, X)$ is $K(\mathbf{x}_i, \mathbf{x}_j)$.

This is called the **kernel trick**.

Prediction: compute the predictive distribution of y' conditioned on Y :

$$y'| \mathbf{x}', X, Y, K \sim \mathcal{N}(\underbrace{K(\mathbf{x}', X)K(X, X)^{-1} Y^T}_{\text{mean}}, \underbrace{K(\mathbf{x}', \mathbf{x}') - K(\mathbf{x}', X)K(X, X)^{-1}K(X, \mathbf{x}')}_{\text{variance}})$$

Regression using the covariance kernel

For non-linear regression, all operations depended on $K(\mathbf{x}, \mathbf{x}')$ rather than explicitly on $\phi(\mathbf{x})$.

So we can define the joint in terms of K *implicitly* using a (potentially infinite-dimensional) feature map $\phi(\mathbf{x})$.

$$Y|X, K \sim \mathcal{N}(0_N, K(X, X))$$

where the i, j entry in the covariance matrix $K(X, X)$ is $K(\mathbf{x}_i, \mathbf{x}_j)$.

This is called the **kernel trick**.

Prediction: compute the predictive distribution of y' conditioned on Y :

$$y'| \mathbf{x}', X, Y, K \sim \mathcal{N}(\underbrace{K(\mathbf{x}', X)K(X, X)^{-1}Y^T}_{\text{mean}}, \underbrace{K(\mathbf{x}', \mathbf{x}') - K(\mathbf{x}', X)K(X, X)^{-1}K(X, \mathbf{x}')}_{\text{variance}})$$

Evidence: this is just the Gaussian likelihood:

$$P(Y|X, K) = |2\pi K(X, X)|^{-\frac{1}{2}} e^{-\frac{1}{2} Y K(X, X)^{-1} Y^T}$$

Regression using the covariance kernel

For non-linear regression, all operations depended on $K(\mathbf{x}, \mathbf{x}')$ rather than explicitly on $\phi(\mathbf{x})$.

So we can define the joint in terms of K *implicitly* using a (potentially infinite-dimensional) feature map $\phi(\mathbf{x})$.

$$Y|X, K \sim \mathcal{N}(0_N, K(X, X))$$

where the i, j entry in the covariance matrix $K(X, X)$ is $K(\mathbf{x}_i, \mathbf{x}_j)$.

This is called the **kernel trick**.

Prediction: compute the predictive distribution of y' conditioned on Y :

$$y'| \mathbf{x}', X, Y, K \sim \mathcal{N}(\underbrace{K(\mathbf{x}', X)K(X, X)^{-1}Y^T}_{\text{mean}}, \underbrace{K(\mathbf{x}', \mathbf{x}') - K(\mathbf{x}', X)K(X, X)^{-1}K(X, \mathbf{x}')}_{\text{variance}})$$

Evidence: this is just the Gaussian likelihood:

$$P(Y|X, K) = |2\pi K(X, X)|^{-\frac{1}{2}} e^{-\frac{1}{2} YK(X, X)^{-1} Y^T}$$

Evidence optimisation: the covariance kernel K often has parameters, and these can be optimized by gradient ascent in $\log P(Y|X, K)$.

The Gaussian process

A **Gaussian process** (GP) is a collection of random variables, any finite number of which have (consistent) Gaussian distributions.

In our regression setting, for each input vector \mathbf{x} we have an output $f(\mathbf{x})$. Given $X = [\mathbf{x}_1, \dots, \mathbf{x}_N]$, the joint distribution of the outputs $F = [f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)]$ is:

$$F|X, K \sim \mathcal{N}(0, K(X, X))$$

Thus the random function $f(\mathbf{x})$ (as a collection of random variables, one $f(\mathbf{x})$ for each \mathbf{x}) is a Gaussian process.

The Gaussian process

A **Gaussian process** (GP) is a collection of random variables, any finite number of which have (consistent) Gaussian distributions.

In our regression setting, for each input vector \mathbf{x} we have an output $f(\mathbf{x})$. Given $X = [\mathbf{x}_1, \dots, \mathbf{x}_N]$, the joint distribution of the outputs $F = [f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)]$ is:

$$F|X, K \sim \mathcal{N}(0, K(X, X))$$

Thus the random function $f(\mathbf{x})$ (as a collection of random variables, one $f(\mathbf{x})$ for each \mathbf{x}) is a Gaussian process.

In general, a Gaussian process is parametrized by a **mean function** $m(\mathbf{x})$ and **covariance kernel** $K(\mathbf{x}, \mathbf{x}')$, and we write

$$f(\cdot) \sim \mathcal{GP}(m(\cdot), K(\cdot, \cdot))$$

The Gaussian process

A **Gaussian process** (GP) is a collection of random variables, any finite number of which have (consistent) Gaussian distributions.

In our regression setting, for each input vector \mathbf{x} we have an output $f(\mathbf{x})$. Given $X = [\mathbf{x}_1, \dots, \mathbf{x}_N]$, the joint distribution of the outputs $F = [f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)]$ is:

$$F|X, K \sim \mathcal{N}(0, K(X, X))$$

Thus the random function $f(\mathbf{x})$ (as a collection of random variables, one $f(\mathbf{x})$ for each \mathbf{x}) is a Gaussian process.

In general, a Gaussian process is parametrized by a **mean function** $m(\mathbf{x})$ and **covariance kernel** $K(\mathbf{x}, \mathbf{x}')$, and we write

$$f(\cdot) \sim \mathcal{GP}(m(\cdot), K(\cdot, \cdot))$$

Posterior Gaussian process: on observing X and F , the conditional joint distribution of $F' = [f(\mathbf{x}'_1), \dots, f(\mathbf{x}'_M)]$ on another set of input vectors $\mathbf{x}'_1, \dots, \mathbf{x}'_M$ is still Gaussian:

$$F'|X', X, F, K \sim \mathcal{N}(K(X', X)K(X, X)^{-1}F^\top, K(X', X') - K(X', X)K(X, X)^{-1}K(X, X'))$$

thus the posterior over functions $f(\cdot)|X, F$ is still a Gaussian process!

Regression with Gaussian processes

We seek to learn the function that maps inputs $\mathbf{x}_1, \dots, \mathbf{x}_N$ to outputs y_1, \dots, y_N .

Regression with Gaussian processes

We seek to learn the function that maps inputs $\mathbf{x}_1, \dots, \mathbf{x}_N$ to outputs y_1, \dots, y_N .

Instead of assuming a specific form, we consider a random function drawn from a GP prior:

$$f(\cdot) \sim \mathcal{GP}(0, K(\cdot, \cdot)).$$

Any function is possible (no restriction on support) but some are (much) more likely *a priori*.

Regression with Gaussian processes

We seek to learn the function that maps inputs $\mathbf{x}_1, \dots, \mathbf{x}_N$ to outputs y_1, \dots, y_N .

Instead of assuming a specific form, we consider a random function drawn from a GP prior:

$$f(\cdot) \sim \mathcal{GP}(0, K(\cdot, \cdot)).$$

Any function is possible (no restriction on support) but some are (much) more likely *a priori*.

Observations y_i usually taken to be noisy versions of latent $f(\mathbf{x}_i)$:

$$y_i | \mathbf{x}_i, f(\cdot) \sim \mathcal{N}(f(\mathbf{x}_i), \sigma^2)$$

Regression with Gaussian processes

We seek to learn the function that maps inputs $\mathbf{x}_1, \dots, \mathbf{x}_N$ to outputs y_1, \dots, y_N .

Instead of assuming a specific form, we consider a random function drawn from a GP prior:

$$f(\cdot) \sim \mathcal{GP}(0, K(\cdot, \cdot)).$$

Any function is possible (no restriction on support) but some are (much) more likely *a priori*.

Observations y_i usually taken to be noisy versions of latent $f(\mathbf{x}_i)$:

$$y_i | \mathbf{x}_i, f(\cdot) \sim \mathcal{N}(f(\mathbf{x}_i), \sigma^2)$$

Evidence: given by the multivariate Gaussian likelihood:

$$P(Y|X) = |2\pi(K(X, X) + \sigma^2 I)|^{-\frac{1}{2}} e^{-\frac{1}{2} Y(K(X, X) + \sigma^2 I)^{-1} Y^T}$$

Regression with Gaussian processes

We seek to learn the function that maps inputs $\mathbf{x}_1, \dots, \mathbf{x}_N$ to outputs y_1, \dots, y_N .

Instead of assuming a specific form, we consider a random function drawn from a GP prior:

$$f(\cdot) \sim \mathcal{GP}(0, K(\cdot, \cdot)).$$

Any function is possible (no restriction on support) but some are (much) more likely *a priori*.

Observations y_i usually taken to be noisy versions of latent $f(\mathbf{x}_i)$:

$$y_i | \mathbf{x}_i, f(\cdot) \sim \mathcal{N}(f(\mathbf{x}_i), \sigma^2)$$

Evidence: given by the multivariate Gaussian likelihood:

$$P(Y|X) = |2\pi(K(X, X) + \sigma^2 I)|^{-\frac{1}{2}} e^{-\frac{1}{2} Y(K(X, X) + \sigma^2 I)^{-1} Y^T}$$

Posterior: also a GP:

$$f(\cdot) | X, Y \sim \mathcal{GP}(K(\cdot, X)(K(X, X) + \sigma^2 I)^{-1} Y^T, K(\cdot, \cdot) - K(\cdot, X)(K(X, X) + \sigma^2 I)^{-1} K(X, \cdot))$$

Regression with Gaussian processes

We seek to learn the function that maps inputs $\mathbf{x}_1, \dots, \mathbf{x}_N$ to outputs y_1, \dots, y_N .

Instead of assuming a specific form, we consider a random function drawn from a GP prior:

$$f(\cdot) \sim \mathcal{GP}(0, K(\cdot, \cdot)).$$

Any function is possible (no restriction on support) but some are (much) more likely *a priori*.

Observations y_i usually taken to be noisy versions of latent $f(\mathbf{x}_i)$:

$$y_i | \mathbf{x}_i, f(\cdot) \sim \mathcal{N}(f(\mathbf{x}_i), \sigma^2)$$

Evidence: given by the multivariate Gaussian likelihood:

$$P(Y|X) = |2\pi(K(X, X) + \sigma^2 I)|^{-\frac{1}{2}} e^{-\frac{1}{2} Y(K(X, X) + \sigma^2 I)^{-1} Y^T}$$

Posterior: also a GP:

$$f(\cdot) | X, Y \sim \mathcal{GP}(K(\cdot, X)(K(X, X) + \sigma^2 I)^{-1} Y^T, K(\cdot, \cdot) - K(\cdot, X)(K(X, X) + \sigma^2 I)^{-1} K(X, \cdot))$$

Predictions: posterior on f , plus observation noise:

$$y' | X, Y, \mathbf{x}' \sim \mathcal{N}(E[f(\mathbf{x}') | X, Y], \text{Var}[f(\mathbf{x}') | X, Y] + \sigma^2)$$

Regression with Gaussian processes

We seek to learn the function that maps inputs $\mathbf{x}_1, \dots, \mathbf{x}_N$ to outputs y_1, \dots, y_N .

Instead of assuming a specific form, we consider a random function drawn from a GP prior:

$$f(\cdot) \sim \mathcal{GP}(0, K(\cdot, \cdot)).$$

Any function is possible (no restriction on support) but some are (much) more likely *a priori*.

Observations y_i usually taken to be noisy versions of latent $f(\mathbf{x}_i)$:

$$y_i | \mathbf{x}_i, f(\cdot) \sim \mathcal{N}(f(\mathbf{x}_i), \sigma^2)$$

Evidence: given by the multivariate Gaussian likelihood:

$$P(Y|X) = |2\pi(K(X, X) + \sigma^2 I)|^{-\frac{1}{2}} e^{-\frac{1}{2} Y(K(X, X) + \sigma^2 I)^{-1} Y^T}$$

Posterior: also a GP:

$$f(\cdot) | X, Y \sim \mathcal{GP}(K(\cdot, X)(K(X, X) + \sigma^2 I)^{-1} Y^T, K(\cdot, \cdot) - K(\cdot, X)(K(X, X) + \sigma^2 I)^{-1} K(X, \cdot))$$

Predictions: posterior on f , plus observation noise:

$$y' | X, Y, \mathbf{x}' \sim \mathcal{N}(E[f(\mathbf{x}') | X, Y], \text{Var}[f(\mathbf{x}') | X, Y] + \sigma^2)$$

Evidence Optimisation: gradient ascent in $\log P(Y|X)$.



Samples from a Gaussian process

We can draw sample functions from a GP by fixing a set of input vectors $\mathbf{x}_1, \dots, \mathbf{x}_N$, and drawing a sample $f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)$ from the corresponding multivariate Gaussian. This can then be plotted.

Samples from a Gaussian process

We can draw sample functions from a GP by fixing a set of input vectors $\mathbf{x}_1, \dots, \mathbf{x}_N$, and drawing a sample $f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)$ from the corresponding multivariate Gaussian. This can then be plotted.



Example prior and posterior GPs:



Samples from a Gaussian process

We can draw sample functions from a GP by fixing a set of input vectors $\mathbf{x}_1, \dots, \mathbf{x}_N$, and drawing a sample $f(\mathbf{x}_1), \dots, f(\mathbf{x}_N)$ from the corresponding multivariate Gaussian. This can then be plotted.


Example prior and posterior GPs:



Another approach is to

- ▶ sample $f(\mathbf{x}_1)$ first,
- ▶ then $f(\mathbf{x}_2)|f(\mathbf{x}_1)$,
- ▶ and generally $f(\mathbf{x}_n)|f(\mathbf{x}_1), \dots, f(\mathbf{x}_{n-1})$ for $n = 1, 2, \dots$


Sample from a 2D Gaussian process



Examples of covariance kernels


- Polynomial:

$$K(\mathbf{x}, \mathbf{x}') = (1 + \mathbf{x}^T \mathbf{x}')^m \quad m = 1, 2, \dots$$




- Squared-exponential (or exponentiated-quadratic):

$$K(\mathbf{x}, \mathbf{x}') = \theta^2 e^{-\frac{\|\mathbf{x} - \mathbf{x}'\|^2}{2\eta^2}}$$




- Periodic (exp-sine):

$$K(x, x') = \theta^2 e^{-\frac{2 \sin^2(\pi(x-x')/\tau)}{\eta^2}}$$



- Rational Quadratic:

$$K(\mathbf{x}, \mathbf{x}') = \left(1 + \frac{\|\mathbf{x} - \mathbf{x}'\|^2}{2\alpha\eta^2}\right)^{-\alpha} \quad \alpha > 0$$



Forms of kernels

If K_1 and K_2 are covariance kernels, then so are:

- ▶ Rescaling: αK_1 for $\alpha > 0$.
- ▶ Addition: $K_1 + K_2$
- ▶ Elementwise product: $K_1 K_2$
- ▶ Mapping: $K_1(\phi(\mathbf{x}), \phi(\mathbf{x}'))$ for some function ϕ .

Forms of kernels

If K_1 and K_2 are covariance kernels, then so are:

- ▶ Rescaling: αK_1 for $\alpha > 0$.
- ▶ Addition: $K_1 + K_2$
- ▶ Elementwise product: $K_1 K_2$
- ▶ Mapping: $K_1(\phi(\mathbf{x}), \phi(\mathbf{x}'))$ for some function ϕ .

A covariance kernel is **translation-invariant** if

$$K(\mathbf{x}, \mathbf{x}') = h(\mathbf{x} - \mathbf{x}')$$

A GP with a translation-invariant covariance kernel is stationary: if $f(\cdot) \sim \mathcal{GP}(0, K)$, then so is $f(\cdot - \mathbf{x}) \sim \mathcal{GP}(0, K)$ for each \mathbf{x} .

Forms of kernels

If K_1 and K_2 are covariance kernels, then so are:

- ▶ Rescaling: αK_1 for $\alpha > 0$.
- ▶ Addition: $K_1 + K_2$
- ▶ Elementwise product: $K_1 K_2$
- ▶ Mapping: $K_1(\phi(\mathbf{x}), \phi(\mathbf{x}'))$ for some function ϕ .

A covariance kernel is **translation-invariant** if

$$K(\mathbf{x}, \mathbf{x}') = h(\mathbf{x} - \mathbf{x}')$$

A GP with a translation-invariant covariance kernel is stationary: if $f(\cdot) \sim \mathcal{GP}(0, K)$, then so is $f(\cdot - \mathbf{x}) \sim \mathcal{GP}(0, K)$ for each \mathbf{x} .

A covariance kernel is **radial** or **radially symmetric** if

$$K(\mathbf{x}, \mathbf{x}') = h(\|\mathbf{x} - \mathbf{x}'\|)$$

A GP with a radial covariance kernel is stationary with respect to translations, rotations, and reflections of the input space.

Nonparametric Bayesian Models and Occam's Razor

Overparameterised models can **overfit**. In the GP, the parameter is the function $f(\mathbf{x})$ which can be infinite-dimensional.

Nonparametric Bayesian Models and Occam's Razor

Overparameterised models can **overfit**. In the GP, the parameter is the function $f(\mathbf{x})$ which can be infinite-dimensional.

However, the Bayesian treatment integrates over these parameters: we never identify a single “best fit” f , just a posterior (and posterior mean). So f cannot be adjusted to overfit the data.

Nonparametric Bayesian Models and Occam's Razor

Overparameterised models can **overfit**. In the GP, the parameter is the function $f(\mathbf{x})$ which can be infinite-dimensional.

However, the Bayesian treatment integrates over these parameters: we never identify a single “best fit” f , just a posterior (and posterior mean). So f cannot be adjusted to overfit the data.

The GP is an example of the larger class of **nonparametric Bayesian models**.

- ▶ Infinite number of parameters.
- ▶ Often constructed as the infinite limit of a nested family of finite models (sometimes equivalent to infinite model averaging).
- ▶ Parameters integrated out, so effective number of parameters to overfit is zero or small (hyperparameters).
- ▶ No need for model selection. Bayesian posterior on parameters will concentrate on “submodel” with largest integral automatically.
- ▶ No explicit need for Occam's razor, validation or added regularisation penalty.